



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A





NRL Memorandum Report 5305

Early Time HANE Structure

J. D. HUBA

Geophysical and Plasma Dynamics Branch
Plasma Physics Division

April 30, 1984

This research was sponsored by the Defense Nuclear Agency under Subtask S99QMXBC, work unit 00119 and work unit title "Early Time Dynamics."







NAVAL RESEARCH LABORATORY Washington, D.C.

Approved for public release; distribution unlimited.

84 05 21 035

SECURITY.	ASSIFICATION	· . F	-HIS 6	PAGE
-----------	--------------	-------	--------	------

	REPORT DOCUM	ENTATION PAG	F			
A REPORT SECURITY CLASSIF CATION UNCLASSIFIED		15 RESTRICTIVE MARKINGS				
24 SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION AVAILABILITY OF REPORT				
26 DECLASSIFICATION COWNGRADING SCHEDULE		Approved for public release; distribution unlimited.			n unlimited.	
4 PERFORMING ORGANIZATION REPORT NUMBERIS)		5. MONITORING ORGANIZATION REPORT NUMBERISI				
NRL Memorandum Report 5305		1				
6a NAME OF PERFORMING ORGANIZATION Naval Research Laboratory	56 OFFICE SYMBOL If applicable	78. NAME OF MONITORING ORGANIZATION				
6c ADDRESS (City State and /IP Code)	<u></u>		76 ADDRESS (City State and /IP Code)			÷ ,, ,
Washington, DC 20375						. •
84. NAME OF FUNDING SPONSORING ORGANIZATION	86 OFFICE SYMBOL (If applicable)	9. PROCUREMENT	INSTRUMENT ID	ENTIFICATION		2 1 19
Defense Nuclear Agency Se ADDRESS City State and ZIP Code:	<u></u>			····		1 13
Washington, DC 20305		PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT	Δ
11 TiTLE Include Security Classifications (See page ii)	<u></u>	62715H			47-1944-04	•
(See page ii) 12 PERSONAL AUTHORIS: J. D. Huba			!	<u>L</u>		
13a TYPE OF REPORT 13b. TIME Interim FROM	COVERED	April 30, 1		15 PAGE 29		
16. SUPPLEMENTARY NOTATION Thi						
Subtask S99QMXBC, work unit 00	119 and work unit 18 SUBJECT TERMS A Microturbulence Early time struct	Continue on reverse if no	e Dynamics.'	fy by block numb tability	•••	
Subtask S99QMXBC, work unit 00:	119 and work unit 18 SUBJECT TERMS Microturbulence Early time struct 19 can have a direct ayleigh-Taylor instants ous electron-ion and is developed. It is ansferring energy from the damped modern to the damped modern.	Rayleighture Nonline effect on the evolution collision demonstrated the om the growing e analysis is that	e Dynamics.' ressery and identify gh-Taylor inside ar mode couloution of intred which incoms. A set of at the interm modes to the anomalous eless of this thee	erchange instruction two-dimensine damped molectron-ion cory to early	tabilities. e effects ional length regime odes because ollisions	・ハーラル・フル・ファー・ファー・ファー・ファー・ファー・ファー・ファー・ファー・ファー・ファー
Subtask S99QMXBC, work unit 00. 17 COSATI CODES FIELD GROUP SUB GR 19 ASSTRACT (Continue on reverse if necessery of the sample, two-fluid model of the R of microturbulence through anomal nonlinear mode coupling equations of the instability can saturate by trof the nonlinear E X B motion. A provide the dissipation mechanism is structuring are discussed.	119 and work unit 18 SUBJECT TERMS Microturbulence Early time struct 19 can have a direct ayleigh-Taylor instants ous electron-ion and is developed. It is ansferring energy from the damped modern to the damped modern.	Rayleighture Nonline N	e Dynamics.' ressery and identify gh-Taylor insignar mode con red which income. A set of at the interm modes to the anomalous eless of this theel URITY CLASSIFIE ED	erchange instruction two-dimensine damped molectron-ion cory to early	tabilities. e effects ional length regime odes because ollisions time HANE	

SECURITY CLASSIFICATION OF THIS PAGE	
11. TITLE (Include Security Classification)	
INFLUENCE OF MICROTURBULENCE ON EARLY TIME HANE STRUCTURE	
į daras ir salas sa	
,	

CONTENTS

I.	INT	RODUCTION	1
II.	DER	RIVATION OF MODE COUPLING EQUATIONS	2
III.	ANA	LYSIS OF MODE COUPLING EQUATIONS	6
	A.	Linear Theory	6
		1. Collisionless plasma $(v_{ei} = v_{ii} = 0)$	7
		2. Collisional plasma $(v_{ei} \neq 0 \text{ and } v_{ii} \neq 0) \dots$	7
	В.	Nonlinear Theory	8
IV.	DIS	CUSSION	12
Ackn	owle	dgments	14
Refe	renc	es	15



Accession For	
NTIS GRA&I DTIC TAB Unramownood Justification	000
Re Distribute on/	
Aveilenditty	::(√o*
All prose	angen en e

INFLUENCE OF MICROTURBULENCE ON EARLY TIME HANE STRUCTURE

INTRODUCTION

An important issue to the Defense Nuclear Agency is understanding the physical processes involved in the structuring of the ionosphere following a HANE. The large-scale, long-lasting ionization irregularities produced by a HANE can adversely impact radar and communications systems (i.e., scintillations). Thus, it is crucial to understand the structuring mechanisms associated with HANES in order to obtain a predictive capability which could aid the operation of these systems in a nuclear environment. Recently, there has been renewed interest in the DNA community concerning early time HANE processes (t \(\left\) few sec). For example, DNA is presently funding a major laser experiment at NRL to explore the physics associated with early time HANE evolution.

Among the various early time processes to be understood, the occurrence of plasma microinstabilities and of structure producing processes (i.e., interchange instabilities) are two important areas. To our knowledge, these two areas have been treated somewhat independently of each other. The studies of microturbulence have been directed at explaining the coupling of debris and air plasmas (i.e., momentum exchange), and on the heating of the plasmas (i.e., energy exchange) (Lampe et al., 1975). On the other hand, the studies of early time structuring processes, which have been limited to date, do not directly include microturbulence effects. It should be noted that microturbulent processes have been included indirectly in an early time structure analysis. The work of Brecht et al. (1982) addresses the growth of Rayleigh-Taylor instabilities based upon the conditions prescribed by the microphysics (e.g., thickness, temperature, density within the coupling shell).

The purpose of this report is to suggest a direct effect of microturbulence on the evolution of early time interchange instabilities. Namely, that microinstabilities can provide a dissipation mechanism (via the anomalous transport properties associated with them) which can cause saturation of an interchange instability through a nonlinear mode coupling process. In order to demonstrate this point, we consider a simplified model. We consider a two-fluid plasma model (ions and electrons) which Manuscript approved January 24, 1984.

incorporates a gravitational field (g) and a density inhomogeneity (∇n) . We develop a set of nonlinear mode coupling equations for this situation and show that the intermediate regime of the instability can saturate by transferring energy from the growing modes to the damped modes because of the nonlinear electron $E \times B$ motion. The crucial point in the analysis is that anomalous electron—ion collisions provide the dissipation mechanism for the damped modes.

The organization of the paper is as follows. In the next section we derive the nonlinear mode equations. In Section III we present the results for a two mode coupling process. Finally, in Section IV we discuss the implications of our theory for early time HANE processes and describe our future efforts in this area.

II. DERIVATION OF MODE COUPLING EQUATIONS

The geometry and plasma configuration used in the analysis are shown in Fig. 1. We take the ambient magnetic field to be in the z direction $(\mathfrak{B}=\mathbb{B} \ \hat{\mathbf{e}}_z)$, and the density to be inhomogeneous in the x direction $(\mathfrak{n}=\mathfrak{n}(x))$. The fundamental assumptions used in the analysis are the following. We assume perturbed quantities to be proportional to $\exp[i(k_x x + k_y y - \omega t)]$ with $k_x L >> 1$, $k_y L >> 1$, $k_x \rho_i << 1$, $k_y \rho_i << 1$, and $\omega << \Omega_i$ where $L = (\partial \ln n/\partial x)^{-1}$ is the density gradient scale length $\rho_i = (T_i/m_i)^{1/2}/\Omega_i$ is the mean ion Larmor radius, and $\Omega_i = \mathrm{eB/m_i} c$ is the ion cyclotron frequency. We consider a weakly collisional plasma such that $v_{ei}/\Omega_e << 1$, $v_{ie}/\Omega_i << 1$, and $v_{ii}/\Omega_i << 1$ where $v_{\alpha\beta}$ represents a collision frequency between the α and β species, and $\Omega_e = \mathrm{eB/m_e} c$ is the electron cyclotron frequency. Here, the collisions are attributed to wave-particle interactions, i.e., anomalous collisions. Finally, we assume quasineutrality $(n_e = n_i)$.

The equations used in the analysis are continuity, momentum transfer, and charge conservation:

$$\frac{\partial \mathbf{n}}{\partial \mathbf{t}} + \nabla \cdot (\mathbf{n} \underline{\mathbf{v}}_{\mathbf{q}}) = 0 \tag{1}$$

$$0 = -\frac{e}{m_e} \left(-\nabla \phi + \frac{1}{c} \underbrace{\nabla}_e \times \underbrace{B} \right) - v_e^2 \frac{\nabla n}{n} - v_{ei} (\underbrace{\nabla}_e - \underbrace{\nabla}_i)$$
 (2)

$$\frac{d\mathbf{v_i}}{d\mathbf{t}} = \frac{\mathbf{e}}{\mathbf{m_i}} \left(-\nabla_{\phi} + \frac{1}{\mathbf{c}} \mathbf{v_i} \times \mathbf{B} \right) + \mathbf{g} - \mathbf{v_i}^2 \frac{\nabla \mathbf{n}}{\mathbf{n}} - \mathbf{v_{ie}} (\mathbf{v_i} - \mathbf{v_e}) - \frac{1}{\mathbf{nm_i}} \nabla \cdot \mathbf{v_i}$$
(3)




Figure 1: Plasma configuration and slab geometry. The magnetic field is in the z-direction ($\tilde{B} = \tilde{B} \cdot \tilde{e}_z$), the density gradient and gravity are in the x-direction ($\nabla n = \partial n/\partial x \cdot \tilde{e}_x$ and $g = g \cdot \tilde{e}_x$), and the wave vector is in the xy plane transverse to B.

$$\nabla \cdot \mathbf{J} = \nabla \cdot \left[\mathbf{n} (\mathbf{y}_{\mathbf{i}} - \mathbf{y}_{\mathbf{p}}) \right] = 0 \tag{4}$$

where α denotes species (e,i), $v_{\alpha} = (T_{\alpha}/m_{\alpha})^{1/2}$ is the thermal velocity, and π_{i} is the ion stress tensor such that (Stix, 1969)

$$\frac{1}{n\mathbf{m_i}} \nabla \cdot \mathbf{\bar{q}_i} = -\frac{1}{2} \Omega_i \rho_i^2 [(\nabla^2 + \frac{\nabla \mathbf{n}}{\mathbf{n}} \cdot \nabla) \mathbf{\bar{y}_i} \times \hat{\mathbf{e}_z} + (\hat{\mathbf{e}_z} \times \frac{\nabla \mathbf{n}}{\mathbf{n}} \cdot \nabla) \mathbf{\bar{y}_i}]$$

$$-0.3 \nu_{ii} \rho_i^2 [(\nabla^2 + \frac{\nabla \mathbf{n}}{\mathbf{n}} \cdot \nabla) \mathbf{\bar{y}_i} - (\hat{\mathbf{e}_z} \times \frac{\nabla \mathbf{n}}{\mathbf{n}} \cdot \nabla) \mathbf{\bar{y}_i} \times \hat{\mathbf{e}_z}]. \tag{5}$$

The first term in Eq. (5) represents finite ion Larmor radius corrections (Roberts and Taylor, 1962), while the second term represents ion viscosity effects (i.e., shear stress). The compressional stress term in Eq. (5) has been neglected since it is smaller by a factor $k\rho_i$ than the first two terms (Stix, 1969).

To lowest order the equilibrium particle drifts are given by (from Eqs. (2) and (3))

$$\mathbf{v}_{e0} = -\mathbf{v}_{de} \hat{\mathbf{e}}_{\mathbf{v}} \tag{6}$$

$$v_{i0} = (v_{di} + v_g)\hat{e}_y$$
 (7)

where $V_{d\alpha} = (v_{\alpha}^2/\Omega_{\alpha})\partial \ln n/\partial x$ is the diamagnetic drift velocity of species α and $V_{\alpha} = -g/\Omega_{\alpha}$ is the ion gravitational drift velocity.

We now perturb Eqs. (1) - (4) about this equilibrium and let n = n₀ + δn , $\phi = \delta \phi$, and $v_{\alpha} = v_{\alpha 0} + \delta v_{\alpha 1} + \delta v_{\alpha 2}$. Here, $\delta v_{\alpha 1}$ is the first order velocity perturbation and $\delta v_{\alpha 2}$ is the second order velocity perturbation (i.e., proportional to small factors such as v_{ei}/Ω_e , v_{ie}/Ω_i , v_{ii}/Ω_i , $k^2 \rho_i^2$, $\Omega_i^{-1} d/dt$). The first order perturbed velocities are

$$\delta \mathbf{v}_{e1} = -\frac{\mathbf{v}_{e}^{2}}{\Omega_{e}} \nabla \left(\frac{e\delta\phi}{T_{e}} - \frac{\delta\mathbf{n}}{\mathbf{n}_{0}}\right) \times \hat{\mathbf{e}}_{z}$$
 (8)

$$\delta \underline{v}_{11} = -\frac{v_1^2}{\Omega_1} \nabla \psi \times \hat{e}_z$$
 (9)

where $\psi = e\delta\phi/T_1 + \delta n/n_0$.

The second order velocities are

$$\delta \mathbf{y}_{e2} = -\frac{\mathbf{v}_{ei}}{\Omega_{e}} \frac{\mathbf{v}_{i}^{2}}{\Omega_{i}} \left(1 + \frac{\mathbf{T}_{e}}{\mathbf{T}_{i}}\right) \nabla \frac{\delta \mathbf{n}}{\mathbf{n}_{0}}$$

$$\delta \mathbf{y}_{i2} = -\rho_{i}^{2} \left(\frac{\partial}{\partial t} + \mathbf{y}_{i0} \cdot \nabla\right) \nabla \psi - \mathbf{v}_{ie} \rho_{i}^{2} \left(1 + \frac{\mathbf{T}_{e}}{\mathbf{T}_{i}}\right) \nabla \frac{\delta \mathbf{n}}{\mathbf{n}_{0}}$$

$$+ \rho_{i}^{2} \frac{1}{2} \frac{\mathbf{v}_{i}^{2}}{\Omega_{i}} \left[\left(\nabla^{2} + \frac{\nabla \mathbf{n}_{0}}{\mathbf{n}_{0}} \cdot \nabla\right) \nabla \psi \times \hat{\mathbf{e}}_{z} + \left(\hat{\mathbf{e}}_{z} \times \frac{\nabla \mathbf{n}_{0}}{\mathbf{n}_{0}} \cdot \nabla\right) \nabla \psi \right]$$

$$+ 0.3 \rho_{i}^{2} \frac{\mathbf{v}_{ii}}{\Omega_{i}} \frac{\mathbf{v}_{i}^{2}}{\Omega_{i}} \left[\left(\nabla^{2} + \frac{\nabla \mathbf{n}_{0}}{\mathbf{n}_{0}} \cdot \nabla\right) \nabla \psi - \left(\hat{\mathbf{e}}_{z} \times \frac{\nabla \mathbf{n}_{0}}{\mathbf{n}_{0}} \cdot \nabla\right) \nabla \psi \times \hat{\mathbf{e}}_{z} \right]$$

$$(11)$$

Substituting Eqs. (8) - (11) into the electron continuity (Eq. (1)) and charge conservation equations (Eq. (4)), we arrive at the coupled set of equations for δn and $\delta \phi$

$$\frac{\partial}{\partial t} \frac{\delta n}{n_0} - \frac{\mathbf{v_i^2}}{\Omega_i} \nabla \frac{e \delta \phi}{T_i} \times \hat{\mathbf{e}}_z \cdot \frac{\nabla n_0}{n_0} - D_e \nabla^2 \frac{\delta n}{n_0} = \frac{\mathbf{v_i^2}}{\Omega_i} \nabla \frac{e \delta \phi}{T_i} \times \hat{\mathbf{e}}_z \cdot \nabla \frac{\delta n}{n_0}$$
(12)

and

$$\frac{\mathbf{g}}{\Omega_{\mathbf{i}}} \times \hat{\mathbf{e}}_{\mathbf{z}} \cdot \nabla \frac{\delta \mathbf{n}}{\mathbf{n}_{\mathbf{0}}} - \rho_{\mathbf{i}}^{2} \left[\left(\frac{\partial}{\partial \mathbf{r}} + \frac{\mathbf{g}}{\Omega_{\mathbf{i}}} \times \hat{\mathbf{e}}_{\mathbf{z}} \cdot \nabla \right) \nabla^{2} - 0.3 \frac{\nabla_{\mathbf{i}}^{2}}{\Omega_{\mathbf{i}}} \frac{\nu_{\mathbf{i}}}{\Omega_{\mathbf{i}}} \nabla^{4} \right] \left(\frac{e\delta\phi}{T_{\mathbf{i}}} + \frac{\delta \mathbf{n}}{\mathbf{n}_{\mathbf{0}}} \right) = 0$$
(13)

where $D_e = v_{ei} \rho_{es}^2$ is the electron diffusion coefficient and $\rho_{es}^2 = (v_i^2/\Omega_e)(1 + T_e/T_i)$.

Thus, Eqs. (12) and (13) represent nonlinear mode coupling equations for the Rayleigh-Taylor instability in the intermediate wavelength regime (i.e., $kL \gg 1$ and $k\rho_i \ll 1$). The physics contained in Eqs. (12) and (13) is described as follows. The second term on the LHS of Eq. (12) leads to growth of the density perturbation while the third term leads to damping because of electron diffusion. The term on the RHS of Eq. (12) is the electron $E \times B$ nonlinearity. The first term on the LHS of Eq. (13) represents the differential motion of the electrons and ions in the y direction because of gravity. The first term in brackets is the ion polarization term while the second term contains ion viscosity effects.

III. ANALYSIS OF MODE COUPLING EQUATIONS

A. Linear Theory

We linearize Eqs. (12) and (13) by neglecting the electron $E \times B$ nonlinearity in Eq. (12). We assume that perturbations are proportional to $\exp[i(k_x x + k_y y - \omega t)]$. From Eq. (12), it can be shown that

$$\frac{\delta n}{n_0} = -\frac{k_y v_i (\rho_i/L)}{\omega + i v_{ei} k^2 \rho_{eg}^2} \frac{e \delta \phi}{T_i}$$
(14)

where L = $(\nabla n_0/n_0)^{-1} = (\partial \ln n_0/\partial x)^{-1}$ is the scale length of the density gradient. Similarly, from Eq. (13) it can be shown that

$$- \left[\omega_{g} + k^{2} \rho_{i}^{2} \left(\omega + \omega_{g} + i \ 0.3 \ v_{ii} k^{2} \rho_{i}^{2} \right) \right] \frac{\delta n}{n_{0}} =$$

$$k^{2} \rho_{i}^{2} \left[\omega + \omega_{g} + i \ 0.3 \ v_{ii} k^{2} \rho_{i}^{2} \right] \frac{e \delta \phi}{T_{i}}$$
(15)

where $\omega_g = k_y g/\Omega_i$. Combining Eqs. (14) and (15), we obtain the linear dispersion equation

$$\omega^2 + b\omega + c = 0 {16}$$

where

$$b = -k_y V_{i0} + i(v_{ei}k^2 \rho_{es}^2 + 0.3 v_{ii}k^2 \rho_i^2)$$
 (17)

$$c = -\frac{k^2}{\frac{y}{k^2}} \frac{g}{L} - \left[k_y \rho_i (\rho_i/L) \Omega_i - i \nu_{ei} k^2 \rho_{es}^2 \right] \times$$
 (18)

$$[-k_y V_g + i \ 0.3 \ v_{ii} k^2 \rho_i^2]$$

and $V_{i0} = V_{di} + V_g$, $V_{di} = v_i^2/\Omega_i L$ and $V_g = -g/\Omega_i$. Equation (16) has the solution

$$\omega = -b/2 + (b^2/4 - c)^{1/2}$$
 (19)

which is rather messy given the complexity of b and c (Eqs. (17) and (18)). In order to gain insight into the solution of Eq. (19), it is best to consider several limiting cases.

1. Collisionless plasma ($v_{ei} = v_{ii} = 0$)

In the collisionless limit, the dispersion equation is given by

$$\omega^{2} - k_{y} V_{10} \omega - \frac{k_{y}^{2}}{k^{2}} \frac{g}{L} = 0$$
 (20)

which is a well known result (Roberts and Taylor, 1962). The solution to Eq. (20) is

$$\omega = \frac{1}{2} k_y V_{i0} + \frac{1}{2} \left[k_y^2 V_{i0}^2 + 4 \frac{k_y^2}{k^2} \frac{g}{L} \right]^{1/2}$$
 (21)

In the limit $k^2V_{10}^2 \ll 4g/L$, the eigenfrequency is

$$\omega = \pm \frac{k_y}{k} \left(\frac{g}{L}\right)^{1/2} \tag{22}$$

so that instability results if g and ∇n are oppositely directed, i.e., g/L < 0. On the other hand, in the limit $k^2 V_{10}^2 > 4g/L$, the mode becomes stable ($\gamma = 0$) and propagating ($\omega_r \sim k_y V_{10}$). Since it is generally the case that $V_{10} \simeq V_{di} = (\rho_i/L)v_i$, the stabilization criterion is roughly

$$k^2 \rho_i^2 > 4gL/v_i^2 \tag{23}$$

and is therefore known as 'finite Larmor radius stabilization'. An important point is that the stable modes have $\gamma = 0$ and are not damped, i.e., $\gamma < 0$. This is a cruetial consideration in the mode coupling saturation mechanism since damped modes are required as energy sinks.

2. Collisional plasma ($v_{ei} \neq 0$ and $v_{ii} \neq 0$)

We now consider the strong collision limit given by $v_{ii}k^2\rho_i^2 >> \omega$, $k_y v_{i0}$. For this case the dispersion equation

$$i\omega \ 0.3 \ v_{ii} k^2 \rho_i^2 = \frac{k^2}{k^2} \frac{g}{L} + 0.3 \ v_{ii} k^2 \rho_i^2 v_{ei} k^2 \rho_{es}^2$$
 (24)

which has the solution

$$\omega = -i \left[\frac{k_y^2}{k^2} \frac{g/L}{0.3 v_{ii} k^2 \rho_i^2} + v_{ei} k^2 \rho_{es}^2 \right]$$
 (25)

Again instability can result for g/L < 0 (oppositely directed g and $\forall n$) as long as

$$\frac{k_y^2}{k^2} \frac{|g/L|}{0.3 \, v_{ij} k^2 \rho_i^2} > v_{ei} k^2 \rho_{es}^2. \tag{26}$$

In the opposite limit,

$$v_{ei}k^2\rho_{es}^2 > \frac{k_y^2}{k^2} \frac{|g/L|}{0.3 v_{ii}k^2\rho_{i}^2}$$
 (27)

the modes are damped, i.e., $\gamma < 0$. The important point is that the electron-ion collision term provides a dissipation mechanism and can produce modes with a negative growth rate. Hence, anomalous electron-ion collisions can produce damped modes in the collisional limit ($\gamma < 0$). This is in contrast to collisionless finite Larmor radius stabilization which produces stable propagating waves ($\gamma = 0$). Finally, we note the similarity of Eq. (25) to that of the usual collisional Rayleigh-Taylor mode which has a growth rate $\gamma = -(k_y^2/k^2)(g/L\nu_{in})$. In the present case, the quantity 0.3 $\nu_{ij}k^2\rho_i^2$ replaces ν_{in} (Hudson and Kennel, 1975).

B. Nonlinear Theory

To illustrate the nonlinear saturation of the Rayleigh-Taylor instability we consider the collisional limit and follow the analysis prescribed in Rognlein and Weinstock (1974). Mathematically, this analysis closely follows that of Chaturvedi and Ossakow (1977), but the underlying physical concepts and applications are substantially different. The equations of interest are

$$\frac{\partial}{\partial t} \frac{\delta n}{n_0} - \frac{\mathbf{v_i^2}}{\Omega_i} \nabla \frac{e \delta \phi}{T_i} \times \hat{\mathbf{e}}_z \cdot \frac{\nabla n_0}{n_0} - \mathbf{p}_e \nabla^2 \frac{\delta n}{n_0} = \frac{\mathbf{v_i^2}}{\Omega_i} \nabla \frac{e \delta \phi}{T_i} \times \hat{\mathbf{e}}_z \cdot \nabla \frac{\delta n}{n_0}$$
 (28)

and

$$\frac{\mathbf{\xi}}{\Omega_{\mathbf{i}}} \times \hat{\mathbf{e}}_{\mathbf{z}} \cdot \nabla \frac{\delta \mathbf{n}}{\mathbf{n}_{0}} + 0.3 \, \nu_{\mathbf{i}\mathbf{i}} \rho_{\mathbf{i}}^{4} \, \nabla^{4} \left(\frac{\mathbf{e}\delta \phi}{\mathbf{T}_{\mathbf{i}}} + \frac{\delta \mathbf{n}}{\mathbf{n}_{0}} \right) = 0$$
 (29)

where Eq. (28) is the same as Eq. (12) and Eq. (29) follows from Eq. (13) by neglecting ion inertial effects.

Equation (28) is linear and one can obtain a simple relationship between $\delta \phi$ and δn based upon it, namely,

$$\frac{e\delta\phi}{T_i} = -i \Gamma \frac{\delta n}{n_0} \tag{30}$$

where

$$\Gamma = -\frac{8}{0.3 v_{ii} k^2 \rho_i^2} \frac{k_y^{\Omega_i}}{k^2 v_i^2}$$
 (31)

and we have assumed $k_y L > \Omega_1/\nu_{11}$.

Equation (27) is nonlinear and can be rewritten as

$$\frac{\partial}{\partial t} \frac{\delta n}{n_0} = \gamma \frac{\delta n}{n_0} + \frac{v_1^2}{\Omega_i} \nabla \frac{e \delta \phi}{T_i} \times \hat{e}_z \cdot \nabla \frac{\delta n}{n_0}$$
 (32)

where γ is the net linear growth (or damping) of modes. The term γ results from a combination of the second term on the LHS of Eq. (28) (the growing term) and the third term on the LHS of Eq. (28) (the damping term), and is given by

$$\gamma = \frac{k_y^2}{k^2} \frac{|g/L|}{0.3 \, v_{t,t} k_{\rho_t}^2} - v_{ei} k_{\rho_{es}}^2$$
 (33)

for g/L < 0.

TREPORTED INTO DESCRIPTION OF THE PROPERTY OF

We substitute a two-dimensional perturbation of the form

$$\frac{\delta n}{n_0} = A_{1,1} \sin(k_y y - \omega t) \cos k_x x \tag{34}$$

and (from Eq. (29))

$$\frac{e\delta\phi}{T_1} = \Gamma A_{1,1} \cos(k_y y - \omega t) \cos k_x x \tag{35}$$

into the nonlinear term of Eq. (32). This procedure yields the following result

$$\frac{\mathbf{v_i^2}}{\Omega_i} \nabla \frac{\mathbf{e}^{\delta \phi}}{T_i} \times \hat{\mathbf{e}}_z \cdot \nabla \frac{\delta \mathbf{n}}{n_0} = \frac{g}{0.6 \, v_{ii} k^2 \rho_i^2} \frac{k_y^2}{k^2} \, k_x A_{1,1} \sin 2 \, k_x x, \tag{36}$$

that is, a second spatial harmonic in the x-direction (along the density gradient) is produced. Thus, we consider a general perturbation of the form

$$\frac{\delta n}{n_0} = A_{1,1} \sin (k_y y - \omega t) \cos k_x x + A_{2,0} \sin 2 k_x x$$
 (37)

Substituting Eq. (37) into Eq. (32) and making use of Eq. (30), we arrive at the following set of coupled nonlinear differential equations for the mode amplitudes

$$\frac{\partial}{\partial t} A_{1,1} = \gamma_{1,1} A_{1,1} - \alpha A_{1,1} A_{2,0}$$
 (38)

and

$$\frac{\partial}{\partial t} A_{2,0} = \gamma_{2,0} A_{2,0} + \frac{\alpha}{2} A_{1,1}^2$$
 (39)

where the coupling coefficient is

$$\alpha = k_{x} \frac{k_{y}^{2}}{k^{2}} \frac{g}{0.3 v_{x,t} k^{2} \rho_{x}^{2}}$$
 (40)

The mode $A_{2,0}$ is linearly damped (i.e., $\gamma_{2,0} < 0$) which is clear from Eq. (33) since $k_y = 0$ for this mode. Thus, the action of the nonlinear E × B term is to transfer energy from a growing mode $(A_{1,1})$ to a damped mode $(A_{2,0})$ via a mode coupling interaction. This is shown schematically in Fig. 2. It is possible then to set up a steady state situation such that $\partial A_{1,1}/\partial t = 0$ and $\partial A_{2,0}/\partial t = 0$. One can show that this results in the following set of steady state amplitudes

$$A_{2,0} = \frac{\gamma_{1,1}}{\alpha} = \frac{1}{k_x L} \tag{41}$$

and

$$A_{1,1} = \left(-\frac{2|\gamma_{2,0}|}{\alpha} A_{2,0}\right)^{1/2}$$

$$= \left(\frac{k^2}{k_v^2} \frac{v_{ei} k^2 \rho_{es}^2 .0.3 v_{ii} k^2 \rho_{i}^2}{gL k_x^2}\right)^{1/2}$$
(42)

We point out that for $|\gamma_{1,1}| > |\gamma_{2,0}|$, one finds that $A_{2,0} > A_{1,1}$ so that the damped modes can be nonlinearily driven to larger amplitudes than the linear driven modes. These results are similar to those of Rognlein and Weinstock (1974), Chaturvedi and Ossakow (1977) and Chaturvedi and Ossakow (1979). Finally, although a rather simple two-mode analysis has been presented, numerical simulations using many modes for both the collisional Rayleigh-Taylor and E × B instability are consistent with this analysis (Keskinen et al., 1980a,b).




Figure 2: Schematic of mode coupling process. The growing and damped modes are shown in k space. The coupling of the growing (1,1) mode and the damped (0,2) mode is also shown.

IV. DISCUSSION

The purpose of this paper has been to suggest an important influence of microturbulence on the early time structuring of a HANE. We consider the question, how do anomalous collisions produced by microturbulence affect the development of interchange instabilities? To answer this question, we have adopted a simple two-fluid model of the intermediate wavelength Rayleigh-Taylor instability as a test problem. Specifically, we have developed a two-dimensional nonlinear mode coupling theory of the Rayleigh-Taylor instability in the presence of microturbulence. primary effect of the microturbulence is to generate anomalous collisions between electrons and ions, as well as ions and ions; an effect considered to be very important in the early time evolution of a HANE. Examples of instabilities investigated for early time HANE applications are the ion-ion counter-streaming instabilities, the modified-two-stream instability, the ion acoustic instability, and the beam cyclotron instability (Lampe et al., 1975). Lampe et al. (1975) have shown that these instabilities can produce significant anomalous collisions between particles, e.g., $\nu_{\mbox{ei}} \ \mbox{\ifootnote \ensuremath{\searrow}} \ \omega_{\mbox{pe}}$ for the modified two stream and $v_{i,i} \lesssim \omega_{i,h}$ for the magnetized ion-ion counterstreaming instabilities where ω_{pe} is the electron plasma frequency and ω_{lh} is the lower hybrid frequency. We note that in the model considered, the concept of anomalous ion-ion collisions for a single ion fluid is unclear, i.e., anomalous collisionless viscosity is not well understood, but it is a valid concept for a multi-ion species plasma. Thus, as noted in the introduction, the problem addressed in this paper should be considered as an idealized test problem which highlights a potentially important effect in early time HANE physics.

The major result of this work is that the intermediate wavelength modes of the Rayleigh-Taylor instability can be stabilized via a mode coupling process in the collisional regime. Basically, wave energy is transferred nonlinearly from growing modes to damped modes, and a steady state can result. Anomalous electron-ion collisions play a crucial role in that they provide a dissipation mechanism which produces a set of damped modes. This is in contrast to stabilization of the instability due to finite Larmor radius effects which does not produce damped modes, but rather propagating modes with $\gamma = 0$.

A considerable amount of work remains to be done on this problem. Within the context of the simple model developed in this paper two avenues of research are presently being pursued. First, numerical analysis is underway to consider the many-mode situation which is more realistic. In this analysis, a pseudo-spectral code is being used which has proved valuable in understanding the mode coupling stabilization of the lower-hybrid-drift instability (Drake et al., 1983a,b). Second, the effects of ion inertia need to be included self-consistently to understand the transition from the collisionless to collisional regime. Again, the numerical techniques being used allow for this.

While the above described work will provide a better understanding of the nonlinear stabilization of the Rayleigh-Taylor instability, it does not directly apply to early time HANE phenomena. It does provide insight into potentially important effects, and the development of relevant numerical tools. However, considerably different models need to be developed for early time HANE behavior. The types of issues that should be addressed are as follows:

- 1. A stability analysis of early time HANE has been performed by Brecht and Papadopoulos (1979). They considered the development of a Rayleigh-Taylor instability driven by forces associated with laminar-like acceleration of ions within the coupling shell and with the curvature of the magnetic field lines. Another effect not considered in Brecht and Papadopoulos (1979) is the deceleration of the coupling shell due to line tying currents to the conducting ionosphere (Pilipp, 1971; Fedder, 1980) which could also lead to a Rayleigh-Taylor instability. The point is that there are several mechanisms that could produce early time structure instabilities and that a nonlinear analysis of these instabilities, incorporating microturbulence via anomalous collisions, is warranted. However, this is somewhat more complicated than the present calculation since at least two ion species are required in the analysis, and electromagnetic effects may need to be included.
- 2. Another possibility for the generation of structure (i.e., density irregularities) in early time HANE is the onset of the Richtmeyer-Meshkov instability (Richtmeyer, 1960; Andronov et al., 1976). This instability grows linearly with time, as opposed to exponentially, and can occur when a

shock wave passes through an irregular (corrugated) interface from a less dense to a more dense fluid. The turbulence generated from this instability may also be influenced by microturbulence, as in the case of the more standard interchange instability case.

3. It has also been suggested that the weapon debris is 'prestructured' by the 'nuts and bolts' of the device itself (Chesnut, private communication). The subsequent evolution of this structure could also be affected by microturbulence. For example, the wavenumber spectrum of the turbulence could be altered by the microturbulence through a mode coupling process.

In conclusion, we have shown that microturbulence, via its associated anomalous collisional effects, can influence the nonlinear development of the collisional Rayleigh-Taylor instability. Namely, it can lead to the stabilization of the intermediate wavelength modes through a two-dimensional mode coupling process. Anomalous electron-ion collisions provide the needed dissipation mechanism. Thus, microturbulence could play an important role in the development of structure for early time HANE phenomena. We have given some examples of the structure problems that could be affected by microturbulence and are relevant to early time HANE processes. Admittedly, some of these ideas are speculative at this time, but it is clear that further research is needed in this area.

Acknowledgments

I wish to thank Drs. P. Chaturvedi, A. Hassam and M. Keskinen for helpful conversations. This work has been supported by the Defense Nuclear Agency under MIPR 83-693.

References

- Andronov, V.A., S.M. Bakrakh, E.E. Meshkov, V.N. Mokhov, V.V. Nikiforov, A.V. Pevnitskii, and A.I. Tolshmyakov, Turbulent mixing at contact surface accelerated by shock waves, Sov. Phys. JETP, 44, 424, 1976.
- Brecht, S.H. and K. Papadopoulos, "Cross Field Jetting of Energetic Ions
 Produced By Raleigh-Taylor Instability," NRL Memo 4068, 1979. (AD-A078 027)
- Brecht, S.H., E. Hyman, and P.J. Palmadesso, Structuring caused by Rayleigh-Taylor instability," NRL Memo 4790, 1982.
- Chaturvedi, P.K., and S.L. Ossakow, Nonlinear theory of the collisional Rayleigh-Taylor instability in equatorial spread F, Geophys. Res. Lett., 4, 558, 1977.
- Chaturvedi, P.K., and S.L. Ossakow, Nonlinear stabilization of the E × B gradient drift instability in ionospheric plasma clouds, J. Geophys. Res., 84, 419, 1979.
- Drake, J.F., P.N. Guzdar, and J.D. Huba, Saturation of the lower-hybrid-drift instability by mode coupling, Phys. Fluids, 26, 601, 1983.
- Drake, J.F., P.N. Guzdar, A.B. Hassam, and J.D. Huba, Nonlinear mode coupling theory of the lower-hybrid-drift instability, submitted to Phys. Fluids, 1983.
- Fedder, J.A., "Structuring of collisionless, high velocity ion clouds," NRL Memo 4307, 1980.
- Hudson, M.K. and C.F. Kennel, Linear theory of equatorial spread F, J. Geophys. Res., 80, 4581, 1975.
- Keskinen, M.J., S.L. Ossakow, and P.K. Chaturvedi, Preliminary report of numerical simulations of intermediate wavelength collisional Rayleigh-Taylor instability in equatorial spread F, J. Geophys. Res., 85, 1775, 1980.
- Keskinen, M.J., S.L. Ossakow, and P.K. Chaturvedi, Preliminary report of numerical simulations of intermediate wavelength E × B gradient drift instability in ionospheric plasma clouds, J. Geophys. Res., 85, 3485, 1980.
- Lampe, M., W.M. Manheimer, and K. Papadopoulos, "Anomalous transport coefficients for HANE applications due to plasma Micro-Instabilities," NRL Memo 3076, 1975.

- Pilipp, W.G., Expansion of an ion cloud in the earth's magnetic field, Planet. Space Sci., 19, 1095, 1971.
- Richtmeyer, R.D., Taylor instability in shock acceleration of compressible fluids, Comm. Pure and Appl. Math., 13, 297, 1960.
- Roberts, K.V. and J.B. Taylor, Magnetohydrodynamic equations for finite Larmor radius, Phys. Rev. Lett., 8, 197, 1962.
- Rognlein, T.D. and J. Weinstock, Theory of the nonlinear spectrum of the gradient drift instability in the equatorial electrojet, J. Geophys. Res., 79, 4733, 1974.
- Stix, T.H., Finite-amplitude collisional drift waves, Phys. Fluids, 12, 627, 1969.

THE PARTY OF THE P

DISTRIBUTION LIST

Andrew Land Company and the Secretary Secretar

DEPARTMENT OF DEFENSE

ASSISTANT SECRETARY OF DEFENSE COMM, CMD, CONT 7 INTELL WASHINGTON, D.C. 20301

DIRECTOR
COMMAND CONTROL TECHNICAL CENTER
PENTAGON RM BE 685
WASHINGTON, D.C. 20301
Olcy ATTN C-650
Olcy ATTN C-312 R. MASON

DIRECTOR
DEFENSE ADVANCED RSCH PROJ AGENCY
ARCHITECT BUILDING
1400 WILSON BLVD.
ARLINGTON, VA. 22209

Olcy ATTN NUCLEAR MONITORING RESEARCH OLCY ATTN STRATEGIC TECH OFFICE

DEFENSE COMMUNICATION ENGINEER CENTER
1860 WIEHLE AVENUE
RESTON, VA. 22090
OICY ATTN CODE R410
OICY ATTN CODE R812

DEFENSE TECHNICAL INFORMATION CENTER CAMERON STATION
ALEXANDRIA, VA. 22314
O2CY

DIRECTOR
DEFENSE NUCLEAR AGENCY
WASHINGTON, D.C. 20305
OICY ATTN STVL
O4CY ATTN TITL
OICY ATTN DDST
O3CY ATTN RAAE

COMMANDER
FIELD COMMAND
DEFENSE NUCLEAR AGENCY
KIRTLAND, AFB, NM 87115
OICY ATTN FCPR

DEFENSE NUCLEAR AGENCY SAO/DNA BUILDING 20676 KIRTLAND AFB, NM 87115 Olcy D.C. THORNBURG

DIRECTOR
INTERSERVICE NUCLEAR WEAPONS SCHOOL
KIRTLAND AFB, NM 87115
01CY ATTN DOCUMENT CONTROL

JOINT CHIEFS OF STAFF
WASHINGTON, D.C. 20301
O1CY ATTN J-3 WWMCCS EVALUATION OFFICE

DIRECTOR
JOINT STRAT TGT PLANNING STAFF
OFFUTT AFB
OMAHA, NB 68113
O1CY ATTN JLTW-2
O1CY ATTN JPST G. GOETZ

CHIEF
LIVERMORE DIVISION FLD COMMAND DNA
DEPARTMENT OF DEFENSE
LAWRENCE LIVERMORE LABORATORY
P.O. BOX 808
LIVERMORE, CA 94550
Olcy Attn FCPRL

COMMANDANT
NATO SCHOOL (SHAPE)
APO NEW YORK 09172
O1CY ATTN U.S. DOCUMENTS OFFICER

UNDER SECY OF DEF FOR RSCH & ENGRG
DEPARTMENT OF DEFENSE
WASHINGTON, D.C. 20301
Olcy ATTN STRATEGIC & SPACE SYSTEMS (OS)

WWMCCS SYSTEM ENGINEERING ORG WASHINGTON, D.C. 20305 Olcy Attn R. CRAWFORD

COMMANDER/DIRECTOR
ATMOSPHERIC SCIENCES LABORATORY
U.S. ARMY ELECTRONICS COMMAND
WHITE SANDS MISSILE RANGE, NM 88002
O1CY ATTN DELAS-EO F. NILES

DIRECTOR

BMD ADVANCED TECH CTR

HUNTSVILLE OFFICE
P.O. BOX 1500

HUNTSVILLE, AL 35807

OLCY ATTN ATC-T MELVIN T. CAPPS
OLCY ATTN ATC-O W. DAVIES
OLCY ATTN ATC-R DON RUSS

PROGRAM MANAGER
BMD PROGRAM OFFICE
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
01CY ATTN DACS-BMT J. SHEA

CHIEF C-E- SERVICES DIVISION
U.S. ARMY COMMUNICATIONS CMD
PENTAGON RM 1B269
WASHINGTON, D.C. 20310
O1CY ATTN C- E-SERVICES DIVISION

COMMANDER
FRADCOM TECHNICAL SUPPORT ACTIVITY
DEPARTMENT OF THE ARMY
FORT MONMOUTH, N.J. 07703
OICY ATTN DRSEL-NL-RD H. BENNET
OICY ATTN DRSEL-PL-ENV H. BOMKE
OICY ATTN J.E. QUIGLEY

COMMANDER
U.S. ARMY COMM-ELEC ENGRG INSTAL AGY
FT. HUACHUCA, AZ 85613
Olcy ATTN CCC-EMEO GEORGE LANE

COMMANDER
U.S. ARMY FOREIGN SCIENCE & TECH CTR
220 7TH STREET, NE
CHARLOTTES VILLE, VA 22901
01CY ATTN DRXST-SD

COMMANDER
U.S. ARMY MATERIAL DEV & READINESS CMD
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
O1CY ATTN DRCLDC J.A. BENDER

COMMANDER
U.S. ARMY NUCLEAR AND CHEMICAL AGENCY
7500 BACKLICK ROAD
BLDG 2073
SPRINGFIELD, VA 22150
O1CY ATTN LIBRARY

DIRECTOR
U.S. ARMY BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MD 21005
O1CY ATTN TECH LIBRARY EDWARD BAICY

COMMANDER
U.S. ARMY SATCOM AGENCY
FT. MONMOUTH, NJ 07703
01CY ATTN DOCUMENT CONTROL

THE RESERVE OF THE PROPERTY OF

COMMANDER
U.S. ARMY MISSILE INTELLIGENCE AGENCY
REDSTONE ARSENAL, AL 35809
Olcy ATTN JIM GAMBLE

DIRECTOR
U.S. ARMY TRADOC SYSTEMS ANALYSIS ACTIVITY
WHITE SANDS MISSILE RANGE, NM 88002
OLCY ATTN ATAA-SA
OLCY ATTN TCC/F. PAYAN JR.
OLCY ATTN ATTA-TAC LTC J. HESSE

COMMANDER
NAVAL ELECTRONIC SYSTEMS COMMAND
WASHINGTON, D.C. 20360
OICY ATTN NAVALEX 034 T. HUGHES
OICY ATTN PME 117
OICY ATTN PME 117-T
OICY ATTN CODE 5011

COMMANDING OFFICER
NAVAL INTELLIGENCE SUPPORT CTR
4301 SUITLAND ROAD, BLDG. 5
WASHINGTON, D.C. 20390
O1CY ATTN MR. DUBBIN STIC 12
O1CY ATTN NISC-50
O1CY ATTN CODE 5404 J. GALET

COMMANDER
NAVAL OCCEAN SYSTEMS CENTER
SAN DIEGO, CA 92152
O1CY ATTN J. FERGUSON

NAVAL RESEARCH LABORATORY
WASHINGTOR, D.C. 20375
OLCY ATTN CODE 4700 S. L. Ossakow
26 GTS IF UNCLASS, 1 CY IF CLASS)
OLCY ATTN CODE 4701 I Vickovitely
OLCY ATTN CODE 4701 I Vickovitely
OLCY ATTN CODE 7500
OLCY ATTN CODE 7500
OLCY ATTN CODE 7500
OLCY ATTN CODE 7500
OLCY ATTN CODE 7551
OLCY ATTN CODE 7551
OLCY ATTN CODE 7555
OLCY ATTN CODE 7555
OLCY ATTN CODE 2755
OLCY ATTN CODE 2028
OLCY ATTN CODE 2028
COMMANDER
NAVAL SEA STEENS COMMAND
WASHINGTOR, D.C. 20362
OLCY ATTN CODE 2228
OLCY ATTN CODE 228
OLCY ATTN CODE 2362
OLCY ATTN CODE 2362
OLCY ATTN CODE 2364
OLCY ATTN CODE 2364
OLCY ATTN CODE 2365
OLCY ATTN CODE 2365
OLCY ATTN CODE 2365
OLCY ATTN CODE 2366
OLCY ATTN CODE 2

WHITE OAK, SILVER SPRING, MD 20910 OICY ATTN CODE F31

DIRECTOR STRATEGIC SYSTEMS PROJECT OFFICE DEPARTMENT OF THE NAVY WASHINGTON, D.C. 20376 Olcy Attn NSP-2141 Olcy Attn NSSP-2722 FRED WIMBERLY

COMMANDER NAVAL SURFACE WEAPONS CENTER DAHLGREN LABORATORY DAHLGREN, VA 22448 OLCY ATTN CODE DF-14 R. BUTLER

OFFICER OF NAVAL RESEARCH ARLINGTON, VA 22217 OLCY ATTN CODE 465 OLCY ATTN CODE 461 OICY ATTN CODE 402 OICY ATTN CODE 420 OLCY ATTN CODE 421

PATRICK AFB, FL 32925 Olcy ATTN TF/MAJ WILEY OLCY ATTN IN

AIR FORCE AVIONICS LABORATORY WRIGHT-PATTERSON AFB, OH 45433 OLCY ATTN AAD WADE HUNT OLCY ATTN AAD ALLEN JOHNSON

DEPUTY CHIEF OF STAFF RESEARCH, DEVELOPMENT, & ACQ DEPARTMENT OF THE AIR FORCE WASHINGTON, D.C. 20330 OLCY ATTN AFRDQ

HEADOUARTERS ELECTRONIC SYSTEMS DIVISION DEPARTMENT OF THE AIR FORCE HANSCOM AFB, MA 01731 OLCY ATTN J. DEAS

HEADOUARTERS ELECTRONIC SYSTEMS DIVISION/YSEA . DEPARTMENT OF THE AIR FORCE HANSCOM AFB, MA 01732 OLCY ATTN YSEA

HEADQUARTERS
ELECTRONIC SYSTEMS DIVISION/DC
DEPARTMENT OF THE AIR FORCE
HANSCOM AFB, MA 01731
01CY ATTN DCKC MAJ J.C. CLARK

COMMANDER
FOREIGN TECHNOLOGY DIVISION, AFSC
WRIGHT-PATTERSON AFB, OH 45433
OICY ATTN NICD LIBRARY
OICY ATTN ETDP B. BALLARD

COMMANDER
ROME AIR DEVELOPMENT CENTER, AFSC
GRIFFISS AFB, NY 13441
Olcy ATTN DOC LIBRARY/TSLD
Olcy ATTN OCSE V. COYNE

SAMSO/SZ POST OFFICE BOX 92960 WORLDWAY POSTAL CENTER LOS ANGELES, CA 90009 (SPACE DEFENSE SYSTEMS) OICY ATTN SZJ

STRATEGIC AIR COMMAND/XPFS
OFFUTT AFB, NB 68113
O1CY ATTN ADWATE MAJ BRUCE BAUER
O1CY ATTN NRT
O1CY ATTN DOK CHIEF SCIENTIST

SAMSO/SK
P.O. BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CA 90009
O1CY ATTN SKA (SPACE COMM SYSTEMS)
M. CLAVIN

SAMSO/MN NORTON AFB, CA 92409 (MINUTEMAN) OLCY ATTN MNNL

COMMANDER
ROME AIR DEVELOPMENT CENTER, AFSC
HANSCOM AFB, MA 01731
01CY ATTN EEP A. LORENTZEN

DEPARTMENT OF ENERGY
LIBRARY ROOM G-042
WASHINGTON, D.C. 20545
Olcy ATTN DOC CON FOR A. LABOWITZ

DEPARTMENT OF ENERGY
ALBUQUERQUE OPERATIONS OFFICE
P.O. BOX 5400
ALBUQUERQUE, NM 87115
O1CY ATTN DOC CON FOR D. SHERWOOD

EG&G, INC.
LOS ALAMOS DIVISION
P.O. BOX 809
LOS ALAMOS, NM 85544
OICY ATTN DOC CON FOR J. BREEDLOVE

UNIVERSITY OF CALIFORNIA
LAWRENCE LIVERMORE LABORATORY
P.O. BOX 808
LIVERMORE, CA 94550
OICY ATTN DOC CON FOR TECH INFO DEPT
OICY ATTN DOC CON FOR L-389 R. OTT
OICY ATTN DOC CON FOR L-31 R. HAGER
OICY ATTN DOC CON FOR L-46 F. SEWARD

LOS ALAMOS NATIONAL LABORATORY
P.O. BOX 1663
LOS ALAMOS, NM 87545
O1CY ATTN DOC CON FOR J. WOLCOTT
O1CY ATTN DOC CON FOR R.F. TASCHEK
O1CY ATTN DOC CON FOR E. JONES
O1CY ATTN DOC CON FOR J. MALIK
O1CY ATTN DOC CON FOR R. JEFFRIES
O1CY ATTN DOC CON FOR J. ZINN
G1CY ATTN DOC CON FOR P. KEATON
O1CY ATTN DOC CON FOR D. WESTERVELT
O1CY ATTN DOC CON FOR D. WESTERVELT

SANDIA LABORATORIES
P.O. BOX 5800

ALBUQUERQUE, NM 87115

O1CY ATTN DOC CON FOR W. BROWN
O1CY ATTN DOC CON FOR A. THORNBROUGH
O1CY ATTN DOC CON FOR T. WRIGHT
O1CY ATTN DOC CON FOR D. DAHLGREN
O1CY ATTN DOC CON FOR 3141
O1CY ATTN DOC CON FOR SPACE PROJECT DIV

SANDIA LABORATORIES
LIVERMORE LABORATORY
P.O. BOX 969
LIVERMORE, CA 94550
OICY ATTN DOC CON FOR B. MURPHEY
OICY ATTN DOC CON FOR T. COOK

OFFICE OF MILITARY APPLICATION DEPARTMENT OF ENERGY WASHINGTON, D.C. 20545 Olcy ATTN DOC CON DR. YO SONG

OTHER GOVERNMENT

DEPARTMENT OF COMMERCE NATIONAL BUREAU OF STANDARDS WASHINGTON, D.C. 20234

OICY (ALL CORRES: ATTN SEC OFFICER FOR)

INSTITUTE FOR TELECOM SCIENCES NATIONAL TELECOMMUNICATIONS & INFO ADMIN BOULDER, CO 80303

OLCY ATTN A. JEAN (UNCLASS ONLY)
OLCY ATTN W. UTLAUT
OLCY ATTN D. CROMBIE
OLCY ATTN L. BERRY

NATIONAL OCEANIC & ATMOSPHERIC ADMIN ENVIRONMENTAL RESEARCH LABORATORIES DEPARTMENT OF COMMERCE BOULDER, CO 80302 Olcy ATTN R. GRUBB OICY ATTN AERONOMY LAB G. REID

DEPARTMENT OF DEFENSE CONTRACTORS

AEROSPACE CORPORATION P.O. BOX 92957 LOS ANGELES, CA 90009 Olcy ATTN I. GARFUNKEL Olcy ATTN T. SALMI Olcy ATTN V. JOSEPHSON Olcy ATTN S. BOWER OLCY ATTN D. OLSEN

ANALYTICAL SYSTEMS ENGINEERING CORP 5 OLD CONCORD ROAD BURLINGTON, MA 01803 OLCY ATTN RADIO SCIENCES

AUSTIN RESEARCH ASSOC., INC. 1901 RUTLAND DRIVE AUSTIN, TX 78758 O1CY ATTN L. SLOAN OLCY ATTN R. THOMPSON

BERKELEY RESEARCH ASSOCIATES, INC. P.O. BOX 983 BERKELEY, CA 94701 OLCY ATTN J. WORKMAN OLCY ATTN C. PRETTIE OLCY ATTN S. BRECHT

BOEING COMPANY, THE P.O. BOX 3707 SEATTLE, WA 98124 Olcy ATTN G. KEISTER
OLCY ATTN D. MURRAY
OLCY ATTN G. HALL
OLCY ATTN J. KENNEY

CHARLES STARK DRAPER LABORATORY, INC. 555 TECHNOLOGY SQUARE CAMBRIDGE, MA 02139 OlCY ATTN D.B. COX OLCY ATTN J.P. GILMORE

COMSAT LABORATORIES LINTHICUM ROAD CLARKSBURG, MD 20734 OLCY ATTN G. HYDE

CORNELL UNIVERSITY DEPARTMENT OF ELECTRICAL ENGINEERING ITHACA, NY 14850 OLCY ATTN D.T. FARLEY, JR.

ELECTROSPACE SYSTEMS, INC. BOX 1359 RICHARDSON, TX 75080 Olcy ATTN H. LOGSTON Olcy ATTN SECURITY (PAUL PHILLIPS)

EOS TECHNOLOGIES, INC. 606 Wilshire Blvd. Santa Monica, Calif 90401 OlCY ATTN C.B. GABBARD

ESL, INC. 495 JAVA DRIVE SUNNY VALE, CA 94086 Olcy ATTN J. ROBERTS Olcy ATTN JAMES MARSHALL

GENERAL ELECTRIC COMPANY SPACE DIVISION VALLEY FORGE SPACE CENTER GODDARD BLVD KING OF PRUSSIA P.O. BOX 8555 PHILADELPHIA, PA 19101 OLCY ATTN M.H. BORTNER SPACE SCI LAB

GENERAL ELECTRIC COMPANY P.O. BOX 1122 SYRACUSE, NY 13201 OLCY ATTN F. REIBERT

GENERAL ELECTRIC TECH SERVICES CO., INC. HMES COURT STREET SYRACUSE, NY 13201 OlCY ATTN G. MILLMAN

GEOPHYSICAL INSTITUTE
UNIVERSITY OF ALASKA
FAIRBANKS, AK 99701
(ALL CLASS ATTN: SECURITY OFFICER)
01CY ATTN T.N. DAVIS (UNCLASS ONLY)
01CY ATTN TECHNICAL LIBRARY
01CY ATTN NEAL BROWN (UNCLASS ONLY)

GTE SYLVANIA, INC.
ELECTRONICS SYSTEMS GRP-EASTERN DIV
77 A STREET
NEEDHAM, MA 02194
01CY ATTN DICK STEINHOF

ESS, INC.
2 ALFRED CIRCLE
BEDFORD, MA 01730
01CY ATTN DONALD HANSEN

ILLINOIS, UNIVERSITY OF 107 COBLE HALL 150 DAVENPORT HOUSE CHAMPAIGN, IL 61820 (ALL CORRES ATTN DAN MCCLELLAND) 01CY ATTN K. YER

INSTITUTE FOR DEFENSE ANALYSES
1801 NO. BEAUREGARD STREET
ALEXANDRIA, VA 22311
OICT ATTN J.M. AEIN
OICY ATTN ERNEST BAUER
OICY ATTN HANS WOLFARD
OICY ATTN JOEL BENGSTON

INTL TEL & TELEGRAPH CORPORATION 500 WASHINGTON AVENUE NUTLEY, NJ 07110 01CY ATTN TECHNICAL LIBRARY

JAYCOR 11011 TORREYANA ROAD P.O. BOX 85154 SAN DIEGO, CA 92138 OICY ATTN J.L. SPERLING JOHNS HOPKINS UNIVERSITY
APPLIED PHYSICS LABORATORY
JOHNS HOPKINS ROAD
LAUREL, MD 20810
OLCY ATTN DOCUMENT LIBRARIAN
OLCY ATTN THOMAS POTEMRA
OLCY ATTN JOHN DASSOULAS

KAMAN SCIENCES CORP P.O. BOX 7463 COLORADO SPRINGS, CO 80933 OlCY ATTN T. MEAGHER

KAMAN TEMPO-CENTER FOR ADVANCED STUDIES 816 STATE STREET (P.O DRAWER QQ) SANTA BARBARA, CA 93102 OLCY ATTN DASIAC OLCY ATTN WARREN S. KNAPP OLCY ATTN WILLIAM MCNAMARA OLCY ATTN B. GAMBILL

LINKABIT CORP 10453 ROSELLE SAN DIEGO, CA 92121 01CY ATTN IRWIN JACOBS

LOCKHEED MISSILES & SPACE CO., INC P.O. BOX 504
SUNNYVALE, CA 94088
OICY ATTN DEPT 60-12
OICY ATTN D.R. CHURCHILL

LOCKHEED MISSILES & SPACE CO., INC.
3251 HANOVER STREET
FALO ALTO, CA 94304
OICY ATTN MARTIN WALT DEPT 52-12
OICY ATTN W.L. IMHOF DEPT 52-12
OICY ATTN RICHARD G. JOHNSON DEPT 52-1:
OICY ATTN J.B. CLADIS DEPT 52-12

MARTIN MARIETTA CORP ORLANDO DIVISION P.O. BOX 5837 ORLANDO, FL 32805 OICY ATTN R. HEFFNER

M.I.T. LINCOLN LABORATORY
P.O. BOX 73
LEXINGTON, MA 02173
OICY ATTN DAVID M. TOWLE
OICY ATTN L. LOUGHLIN
OICY ATTN D. CLARK

MCDONNEL DOUGLAS CORPORATION
5301 BOLSA AVENUE
HUNTINGTON BEACH, CA 92647
OICY ATTN N. HARRIS
OICY ATTN J. MOULE
OICY ATTN GEORGE MROZ
OICY ATTN W. OLSON
OICY ATTN R.W. HALPRIN
OICY ATTN TECHNICAL LIBRARY SERVICES

MISSION RESEARCH CORPORATION
735 STATE STREET
SANTA BARBARA, CA 93101
O1CY ATTN P. FISCHER
O1CY ATTN W.F. CREVIER
O1CY ATTN STEVEN L. GUTSCHE
O1CY ATTN R. BOGUSCH
O1CY ATTN R. HENDRICK
O1CY ATTN RALPH KILB
O1CY ATTN DAVE SOWLE
O1CY ATTN F. FAJEN
O1CY ATTN M. SCHEIBE
O1CY ATTN CONRAD L. LONGMIRE
O1CY ATTN B. WHITE

MISSION RESEARCH CORP.
1720 RANDOLPH ROAD, S.E.
ALBUQUERQUE, NEW MEXICO 87106
OICY R. STELLINGWERF
OICY M. ALME
OICY L. WRIGHT

MITRE CORPORATION, THE
P.O. BOX 208
BEDFORD, MA 01730
O1CY ATTN JOHN MORGANSTERN
O1CY ATTN G. HARDING
O1CY ATTN C.E. CALLAHAN

MITRE CORP
WESTGATE RESEARCH PARK
1820 DOLLY MADISON BLVD
MCLEAN, VA 22101
OICY ATTN W. HALL
OICY ATTN W. FOSTER

PACIFIC-SIERRA RESEARCH CORP 12340 SANTA MONICA BLVD. LOS ANGELES, CA 90025 01CY ATTN E.C. FIELD, JR. PENNSYLVANIA STATE UNIVERSITY
IONOSPHERE RESEARCH LAB
318 ELECTRICAL ENGINEERING EAST
UNIVERSITY PARK, PA 16802
(NO CLASS TO THIS ADDRESS)
OLCY ATTN IONOSPHERIC RESEARCH LAB

PHOTOMETRICS, INC.
4 ARROW DRIVE
WOBURN, MA 01801
01CY ATTN IRVING L. KOFSKY

PHYSICAL DYNAMICS, INC. P.O. BOX 3027 BELLEVUE, WA 98009 O1CY ATTN E.J. FREMOUW

PHYSICAL DYNAMICS, INC. P.O. BOX 10367 OAKLAND, CA 94610 ATTN A. THOMSON

R & D ASSOCIATES
P.O. BOX 9695

MARINA DEL REY, CA 90291
OLCY ATTN FORREST GILMORE
OLCY ATTN WILLIAM B. WRIGHT, JR.
OLCY ATTN ROBERT F. LELEVIER
OLCY ATTN WILLIAM J. KARZAS
OLCY ATTN H. ORY
OLCY ATTN C. MACDONALD
OLCY ATTN R. TURCO
OLCY ATTN L. DERAND
OLCY ATTN U. TSAI

RAND CORPORATION, THE 1700 MAIN STREET SANTA MONICA, CA 90406 01CY ATTN CULLEN CRAIN 01CY ATTN ED BEDROZIAN

RAYTHEON CO. 528 BOSTON POST ROAD SUDBURY, MA 01776 O1CY ATTN BARBARA ADAMS

RIVERSIDE RESEARCH INSTITUTE
330 WEST 42nd STREET
NEW YORK, NY 10036
01CY ATTN VINCE TRAPANI

SCIENCE APPLICATIONS, INC.
1150 PROSPECT PLAZA
LA JOLLA, CA 92037
OICY ATTN LEWIS M. LINSON
OICY ATTN DANIEL A. HAMLIN
OICY ATTN E. FRIEMAN
OICY ATTN E.A. STRAKER
OICY ATTN CURTIS A. SMITH
OICY ATTN JACK MCDOUGALL

SCIENCE APPLICATIONS, INC 1710 GOODRIDGE DR. MCLEAN, VA 22102 ATTN: J. COCKAYNE

SRI INTERNATIONAL

333 RAVENSWOOD AVENUE
MENLO PARK, CA 94025

OlCY ATTN DONALD NEILSON
OLCY ATTN ALAN BURNS
OLCY ATTN G. SMITH
OLCY ATTN R. TSUNODA
OLCY ATTN DAVID A. JOHNSON
OLCY ATTN WALTER G. CHESNUT
OLCY ATTN WALTER JAYE
OLCY ATTN WALTER JAYE
OLCY ATTN TAYL L. LEADABRAND
OLCY ATTN G. CARPENTER
OLCY ATTN G. PRICE
OLCY ATTN R. LIVINGSTON
OLCY ATTN R. LIVINGSTON
OLCY ATTN V. GONZALES
OLCY ATTN D. MCDANIEL

TECHNOLOGY INTERNATIONAL CORP
75 WIGGINS AVENUE
BEDFORD, MA 01730
01CY ATTN W-P. BOQUIST

TOYON RESEARCH CO.
P.O. Box 6890
SANTA BARBARA, CA 93111
Olcy ATTN JOHN ISE, JR.
Olcy ATTN JOEL GARBARINO

TRW DEFENSE & SPACE SYS GROUP
ONE SPACE PARK
REDONDO BEACH, CA 90278
O1CY ATTN R. K. PLEBUCH
O1CY ATTN S. ALTSCHULER
O1CY ATTN D. DEE
O1CY ATTN D. STOCKWELL
SNTF/1575

VISIDYNE
SOUTH BEDFORD STREET
BURLINGTON, MASS 01803
OICY ATTN W. REIDY
OICY ATTN J. CARPENTER
OICY ATTN C. HUMPHREY

的周围(())

.